

THE EFFECTS OF ELEVATED CO₂ ON THE BEHAVIOR AND PHYSIOLOGY OF JUVENILE ROCKFISHES

Fennie, Hamilton W.¹, Scott L. Hamilton¹, Susan M. Sogard², and James P. Barry³

1. Moss Landing Marine Laboratories 2. NOAA's Southwest Fisheries Science Center, National Marine Fisheries Service

3. Monterey Bay Aquarium Research Institute

Introduction: Rapid increases in atmospheric [CO₂] have increased dissolved oceanic [CO₂] (*p*CO₂) and decreased seawater pH (Caldeira and Wickett, 2003). These chemical changes are known as ocean acidification (OA) and have strong negative effects on marine organisms (Kroeker, 2010). Vertebrates are predicted to be less susceptible to changes in *p*CO₂ and pH because they have well developed acid base compensatory mechanisms (Pörtner, 2005); however, in tropical fish species OA negatively affects both behavior and physiology leading to decreased fitness (Munday et al., 2009; Dixon et al., 2010; Ferrari et al., 2011; Allan et al., 2013). Our goals are to understand how OA will affect the behavior and physiology of juvenile rockfishes living in kelp forests within the highly dynamic California Current System (CCS). The CCS is characterized by strong seasonal and diurnal variability in pH due to upwelling and biological processes.

Methods:

- Captured and reared Copper (*Sebastes caurinus*) and Blue Rockfish (*S. mystinus*) for 2-3 months in 4 pH treatments: 8.0, 7.8, 7.5, 7.2
- Measured pH daily and carbonate chemistry weekly
- Tested:
 1. Behavioral lateralization - Right vs. left brain dominance
 2. Critical swimming speed (*U*_{crit}) - Swimming endurance
 3. Aerobic scope – Difference in oxygen consumption between rest and maximum swimming speed




Fig 1. (a) Diver using hand net to catch juvenile rockfishes, (b) double T-maze for testing behavioral lateralization, and (c) juvenile Copper Rockfish in Loligo swim flume during critical swimming speed trial.

Results: OA affects Copper Rockfish behavior and physiology while Blue Rockfish are tolerant to OA

Chemistry Parameter	Control	Mid Century	End of Century	Extreme
Average Daily pH (±SE)	7.92 (±0.006)	7.79 (±0.006)	7.52 (±0.006)	7.33 (±0.009)
Average Weekly DIC (±SE)	2128.4 (±8.621)	2193.6 (±7.706)	2246.5 (±18.233)	2335.6 (±22.626)

Table 1. Measured daily pH and weekly Dissolved Organic Carbon (DIC) ± SE.

1. *S. caurinus* switch from being unbiased to left turn biased as pH decreases while *S. mystinus* do not change bias significantly
2. *S. caurinus* *U*_{crit} decreases with decreasing pH while *S. mystinus* *U*_{crit} is unaffected by changes in pH
3. *S. caurinus* aerobic scope decreases with decreasing pH while *S. mystinus* aerobic scope is unaffected



Sebastes caurinus



Sebastes mystinus




Fig 2. Effects of OA on the behavior and physiology of *S. caurinus* (orange bars) and *S. mystinus* (blue bars). Average relative behavior lateralization for Copper (a) and Blue Rockfish (b), average critical swimming speed (*U*_{crit}) of Copper (c) and Blue Rockfish (d), and average aerobic scope of Copper (e) and Blue Rockfish (f).

Conclusions

- *S. caurinus* have altered behavior and decreased physiological performance as pH decreases
- *S. mystinus* are tolerant to changes in *p*CO₂ and pH
- Species specific responses could lead to differential survival in the face of climate change (Munday et al., 2010; Allan et al., 2013)
- Differences in early life history may explain species specific responses

Acknowledgements: Christian Denney, Emily Donham, Devona Yates, Heather Kramp, Angalee Kirby, Melissa Campana, Chris Lovera
Funding: CA Seagrant, CSU Coast Scholarship, MLML Scholar Award

Allan, B. J. M., Domenici, P., McCormick, M. I., Watson, S.-A., & Munday, P. L. (2013). Elevated CO₂ Affects Predator-Prey Interactions through Altered Performance. *PLoS ONE*, 8(3), e58520. doi:10.1371/journal.pone.0058520. Caldeira, K., & Wickett, M. E. (2003). Anthropogenic carbon and ocean pH. *Nature*, 425(September), 2003. Dixon, D. L., Munday, P. L., & Jones, G. P. (2010). Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecology Letters*, 13(1), 68–75. doi:10.1111/j.1461-0248.2009.01400.x Ferrari, M. C. O., McCormick, M. I., Munday, P. L., Meekan, M. G., Dixon, D. L., Lonnstedt, Ö., & Chivers, D. P. (2011). Putting prey and predator into the CO₂ equation—qualitative and quantitative effects of ocean acidification on predator-prey interactions. *Ecology Letters*, 14(11), 1143–8. doi:10.1111/j.1461-0248.2011.01683.x Kroeker, K. J., Kordas, R. L., Crim, R. N., & Singh, G. G. (2010). Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology Letters*, 13(11), 1419–34. doi:10.1111/j.1461-0248.2010.01518.x Munday, P. L., Dixon, D. L., Donelson, J. M., Jones, G. P., Pratchett, M. S., Devitsina, G. V., & Døving, K. B. (2009). Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proceedings of the National Academy of Sciences of the United States of America*, 106(6), 1848–52. doi:10.1073/pnas.0809996106 Munday, P. L., Dixon, D. L., McCormick, M. I., Meekan, M., Ferrari, M. C. O., & Chivers, D. P. (2010). Replenishment of fish populations is threatened by ocean acidification. *PNAS*, 107(29), 12930–12934. doi:10.1073/pnas.1004519107/-DCSupplemental www.pnas.org/cgi/doi/10.1073/pnas.1004519107 Pörtner, H. O. (2005). Synergistic effects of temperature extremes, hypoxia, and increases in CO₂ on marine animals: From Earth history to global change. *Journal of Geophysical Research*, 110(C9), C09S10. doi:10.1029/2004JC002561